

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
4 April 2002 (04.04.2002)

PCT

(10) International Publication Number  
WO 02/26827 A1(51) International Patent Classification<sup>7</sup>: C07K 14/715,  
14/52, C12N 5/10, A61K 38/00, 38/19(CH). LIAU, Gene [US/US]; 14900 Keley Farm Drive,  
Darnestown, MD 20874 (US).

(21) International Application Number: PCT/EP01/11252

(74) Agent: BECKER, Konrad; Novartis AG, Corporate In-  
tellectual Property, Patent & Trademark Dept., CH-4002  
Basel (CH).(22) International Filing Date:  
28 September 2001 (28.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
1910/00 29 September 2000 (29.09.2000) CH(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,  
ZA, ZW.(71) Applicant (*for all designated States except AT, US*): NO-  
VARTIS AG [CH/CH]; Lichtstrasse 35, CH-4056 Basel  
(CH).(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).(71) Applicant (*for AT only*): NOVARTIS-ERFINDUNGEN  
VERWALTUNGSGESELLSCHAFT M.b.H [AT/AT];  
Brunner Strasse 59, A-1230 Vienna (AT).

## Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): MAR-  
TINY-BARON, Georg [DE/DE]; Am Scheibenberg 10,  
79336 Herbolzheim (DE). WOOD, Jeanette, Marjorie  
[NZ/CH]; In den Kleematten 18, CH-4105 Biel-BenkenFor two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

WO 02/26827 A1

(54) Title: EXTRACELLULAR POLYPEPTIDES OF EPH B RECEPTORS AND EPHRIN B LIGANDS AND THE CORRE-  
SPONDING NUCLEIC ACID MOLECULES

(57) Abstract: The present invention relates to isolated extracellular polypeptides of Eph B receptors and ephrin B ligands and the corresponding isolated nucleic acid molecules comprising the sequences coding for such polypeptides, to recombinant vectors, such as viral vectors, that comprise such nucleic acid molecules, to host cells comprising such recombinant vectors, to pharmaceutical compositions comprising such polypeptides, nucleic acid molecules or recombinant vectors, and to the use of such vectors in the production of pharmaceutical compositions for treating a mammalian disease, including a human disease, especially tumour diseases and ocular neovascularization, by means of gene therapy.

Extracellular polypeptides of Eph B receptors and ephrin B ligands and the corresponding nucleic acid molecules

Field of the Invention:

The present invention relates in particular to isolated extracellular polypeptides of Eph B receptors and ephrin B ligands and the corresponding isolated nucleic acid molecules comprising the sequences coding for such polypeptides, to recombinant vectors, such as viral vectors, that comprise such nucleic acid molecules, to host cells comprising such recombinant vectors, to pharmaceutical compositions comprising such polypeptides, nucleic acid molecules or recombinant vectors, and to the use of such vectors in the production of pharmaceutical compositions for treating a mammalian disease, including a human disease especially tumour diseases and ocular neovascularization, by means of gene therapy.

Background of the Invention:

Eph receptors with 14 different members form the largest sub-group of receptor tyrosine kinases up to now. The extracellular domain consists of a N-terminal globular domain, a cystein-rich domain and two fibronectin type III domains [Himanen *et al.*, Nature 396, 486-491 (1998)].

Owing to the extracellular domain and ligand recognition, the group is sub-divided into two sub-classes: Eph A (A1-A8) and Eph B (B1-B6) [Eph Nomenclature Committee, Cell 90, 403-404 (1997)].

All of the eight ligands known up to now are proteins attached to membrane and are called ephrins. Here also, two sub-classes may be distinguished: ephrin A ligands (A1-A5) are inserted in the membrane by a glycosyl phosphatidyl inositol anchor (GPI), and ephrin B ligands (B1-B3) have a transmembrane domain and a cytoplasmatic protein section consisting of 82-88 amino acids [Eph Nomenclature Committee, Cell 90, 403-404 (1997); Beckmann *et al.*, EMBO J. 13, 3757-3762 (1994)].

Ephrins of the A class bind predominantly to Eph A receptors and ephrins of the B type to Eph B receptors. Within the classes, however, the ligands have overlapping specificities.

Proteins of the Eph/ephrin family play a role in cell and axon orientation. In contrast to chemoattractive substances, Eph/ephrin interactions in the neuronal and epidermal tissues lead to the production of repulsive signals between the cell populations, which express the corresponding receptors and ligands. Using mice whose genes from the Eph/ephrin family have been specifically inactivated, it could be shown that various Eph/ephrins, in particular from the B class, take part in the development of neural crest and in the development of various regions of the brain. Eph/ephrins are thereby made responsible in particular for the complex topography of neuronal systems, which is controlled by direct cell interaction [Brambilla R. and Klein R., *Mol. Cell. Biol.* 15, 4921-4929 (1995); Henkemeyer *et al.*, *Cell* 86, 35-46 (1996)].

Evidence of the participation of the Eph B receptors and ephrin B ligands in embryonal vascular development was provided by genetic experiments. Following homozygote destruction of the ephrin B2 locus, a disturbed formation of the arteries and veins of the capillary vascular system was observed [Wang *et al.*, *Cell* 93, 741-753 (1998)]. In agreement with this, it was shown that ephrin B2 [GeneBank Acc. Nos.: L38847 (murine), U81262 (human); SwissProt Acc. Nos.: P52800 (murine), P52799 (human)] is expressed primarily on arterial endothelial cells, and on the other hand the receptor Eph B4 [GeneBank Acc. Nos.: U06834 (murine), U07695 (human); SwissProt Acc. Nos.: P54761 (murine), P54760 (human)], which binds ephrin B2, is expressed on venous endothelial cells [Wang *et al.*, *Cell* 93, 741-753 (1998)]. To confirm this, targeted inactivation of the Eph B4 gene also exhibits a lethal phenotype similar to ephrin B2, as well as the above-described defects in embryonal vascular development [Gerety *et al.*, *Molecular Cell* 4, 403-414 (1999)]. Further experiments on the function of the Eph B receptor class show, moreover, that Eph B receptors and ephrin B ligands also take part in the cell communication between endothelial cells and the surrounding mesenchymal cells. Mice which cannot express Eph B2 and Eph B3 die *in utero* after about 11.5 days. Analysis of the embryos showed that these animals have defects in the reorganisation of the embryonal vascular system, and exhibited disrupted cell contact between the mesenchymal pericytes and the endodermal endothelial cells [Adams *et al.*, *Genes and Dev.* 13, 295-306 (1999)].

All previous work on the function of Eph B receptors and ephrin B ligands was, however, restricted to the embryonal development of the nervous system and of the embryonal vascular system. Surprisingly, it has now been discovered for the first time that Eph B

receptors and ephrin B ligands in an adult mammal organism participate functionally in angiogenesis. In addition, it can be shown for the first time that Eph B receptors and ephrin B ligands also take part in the pathophysiology of malignant diseases and that extracellular polypeptides of Eph B receptors and ephrin B ligands are suitable for treating tumour diseases. Moreover, it was found that said polypeptides are also suitable for the treatment of ocular neovascularization. It was also surprisingly established that extracellular polypeptides of Eph B receptors and ephrin B ligands possess pharmacokinetic properties of the type which act not only locally but also systemically and have no evident toxic side effects on the organism of mammals.

Definitions:

Within the context of this disclosure the following terms have the meanings as defined below:

In the expression "a soluble polypeptide", the term "soluble" indicates that the respective polypeptide is a non-membrane bound polypeptide which is secreted by mammalian cells which express it. If such a "soluble" polypeptide is expressed for example in cultured mammalian cells, said polypeptide can be found in the cell culture supernatant.

"High affinity" in connection with the binding of ephrin B ligands with Eph B receptors is understood to mean a dissociation constant below 500 nM, preferably in the range of 0.05 nM to 100 nM, more preferably 0.1 nM to 20 nM, the typical range of dissociation constants of ephrin B ligands to Eph B receptors.

The term "essentially similar" used in connection with amino acid sequences refers to amino acid sequence identities. The extracellular regions of different Eph B receptors or ephrin B ligands within the same species have amino acid sequence identities of typically 40% and more among one another. On the other hand, the extracellular regions of the same Eph B receptors or ephrin B ligands of different species typically have amino acid sequence identities of 80% and more. Within the context of this application, a sequence must have at least 40%, preferably at least 75%, most preferably at least 90% amino acid sequence identity to the reference sequence, in order to be regarded as "essentially similar".

Description of the Invention:

- i) The invention relates to an isolated soluble polypeptide which comprises an amino acid sequence of an extracellular region of an Eph B receptor or an essentially similar amino acid sequence thereof, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian ephrin B ligand.
- ii) Similarly the invention also relates to an isolated soluble polypeptide which comprises an amino acid sequence of an extracellular region of an ephrin B ligand or an essentially similar amino acid sequence thereof, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian Eph B receptor.
- iii) The invention further relates especially to an isolated soluble polypeptide which comprises an amino acid sequence of an extracellular region of Eph B4 or an essentially similar amino acid sequence thereof, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian ephrin B ligand.
- iv) Similarly, the invention relates especially to an isolated soluble polypeptide which comprises an amino acid sequence of an extracellular region of ephrin B2 or an essentially similar amino acid sequence thereof, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian Eph B receptor.
- v) An especially preferred embodiment of the invention relates to an isolated soluble polypeptide comprising SEQ ID NO:2 [amino acids (aa) 1-539 of SwissProt Database Accession No. P54761] or SEQ ID NO:4 (aa 1-539 of SwissProt Database Accession No. P54760), which stand for the murine and human extracellular region of Eph B4, respectively, or comprising an amino acid sequence which is essentially similar to SEQ ID NO:2 or SEQ ID NO:4, respectively, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian ephrin B ligand.
- vi) Similarly, a further especially preferred embodiment of the invention relates to an isolated soluble polypeptide which comprises SEQ ID NO:6 (aa 1-229 of SwissProt Database Accession No. P52800) or SEQ ID NO:8 (aa 1-229 of SwissProt Database Accession No. P52799), which stand for the murine and human extracellular region of ephrin B2, respectively, or comprising an amino acid sequence which is essentially similar

to SEQ ID NO:6 or SEQ ID NO:8, respectively, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian Eph B receptor.

vii) The invention also relates to a pharmaceutical composition comprising a polypeptide described in the preceding paragraphs i) – vi), especially such a polypeptide mentioned as being preferred, and to the use of such a polypeptide in particular in the production of a pharmaceutical composition for the treatment of a mammalian, including human disease, especially tumour diseases and ocular neovascularization.

viii) Another preferred embodiment of the invention relates also to an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide described in the preceding paragraphs i) – vi), especially such a polypeptide mentioned as being preferred, to a pharmaceutical composition comprising such a nucleic acid molecule and to the use of such a nucleic acid molecule in particular in the production of a pharmaceutical composition for the treatment of a mammalian, including human disease, especially tumour diseases and ocular neovascularization.

ix) In a further very preferred embodiment the invention relates to a recombinant vector, especially a recombinant viral vector such as a recombinant adenoviral vector, comprising a nucleic acid molecule described in the preceding paragraph viii), especially such a nucleic acid molecule mentioned as being preferred, to a pharmaceutical composition comprising such a recombinant vector and to the use of such a recombinant vector in particular in the production of a pharmaceutical composition for the treatment of a mammalian, including human disease, especially tumour diseases and ocular neovascularization, by means of gene therapy.

x) Another embodiment of the invention relates to a host cell comprising a recombinant vector as described in the preceding paragraph ix), especially such a recombinant vector mentioned as being preferred.

xi) The present invention also relates to a method for the treatment of a tumour disease in a mammal, including human, comprising: effecting an increase, in a mammal in need of such treatment, of the *in vivo* concentration of a polypeptide described in the preceding

paragraphs i) – vi), especially such a polypeptide mentioned as being preferred, to a tumour inhibiting effective amount, wherein said polypeptide has anti-tumour activity *in vivo*.

xii) The present invention further also relates to a method for the treatment of ocular neovascularization in a mammal, including human, comprising: effecting an increase, in a mammal in need of such treatment, of the *in vivo* concentration of a polypeptide described in the preceding paragraphs i) – vi), especially such a polypeptide mentioned as being preferred, to an ocular neovascularization inhibiting effective amount, wherein said polypeptide has anti-ocular neovascularization activity *in vivo*.

The invention most preferably relates to the sequences SEQ ID NO:1-8, especially the human sequences SEQ ID NO:3 [nucleotides (nt) 86-1702 of GeneBank Database Accession No. U07695; nucleotide sequence coding for the extracellular domain of human Eph B4], SEQ ID NO:4, SEQ ID NO:7 (nt 8-694 of GeneBank Database Accession No. U81262; nucleotide sequence coding for the extracellular domain of human ephrin B2) and SEQ ID NO:8, and to their use according to the present invention.

The polypeptides useful in the practice of the present invention (i.e., the polypeptides described in paragraphs i) – vi) above; hereinafter also referred to as “polypeptides of the invention”) can be delivered to an individual in need of treatment using conventional pharmaceutical formulations.

An additional embodiment of the invention relates to the administration of a pharmaceutical composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions contain as active ingredients the polypeptides of the invention or anti-idotypic antibodies to these polypeptides or mimetics of these polypeptides, said active ingredients being administered alone or in combination with at least one other agent, such as a stabilizing agent, in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs or hormones.

The pharmaceutical compositions encompassed by the invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-articular,

intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, Pa.).

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution,



Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc.. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of the polypeptides of the invention, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions where the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

A therapeutically effective dose of active agent can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be

used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example the polypeptides of the invention, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

Normal dosage amounts, e.g., when an exogenously produced polypeptide of the invention is administered, may vary from about 0.1 to about 20 mg/kg per day, preferably from 2.5 to 20 mg/kg per day, depending upon the route and method of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.. Various biodegradable and biocompatible

polymeric matrices, including microcapsules, nanospheres, and implants, are useful in the practice of the present invention.

Microspheres are fine spherical particles containing active drugs. They are differentiated from nanospheres primarily by the size of the particle; microspheres have a diameter of less than approximately 1000  $\mu\text{m}$ , while nanospheres are submicronic ( $<1 \mu\text{m}$ ). Microsphere systems contain either homogeneous monolithic microspheres, in which the drug is dissolved or dispersed homogeneously throughout the polymer matrix, or reservoir-type microspheres, in which the drug is surrounded by the polymer matrix membrane shell. Monolithic and reservoir systems can also be combined. For instance, active drug can be dispersed within, or adsorbed onto, the polymer surface in a reservoir-type microsphere.

Biodegradable polymers can consist of either natural or synthetic materials that vary in purity. Natural polymers include polypeptides and proteins (e.g., albumin, fibrinogen, gelatin, collagen), polysaccharides (e.g., hyaluronic acid, starch, chitosan), and virus envelopes and living cells (e.g., erythrocytes, fibroblasts, myoblasts). Natural materials require cross-linking in the microencapsulation process, leading to the denaturation of the polymer and the embedded drug. As a result, synthetic polymers are most commonly used. Frequently used synthetic polymers include poly(-hydroxy) acids such as polylactic acid (PLA), polyhydroxybutyric acid, and copoly (lactic/glycolic) acid (PLGA). These compounds are biocompatible, lack immunogenicity, and have physical properties that permit them to be easily shaped (to control the bioerosion rate).

Colloidal particulate carriers can also be used in the methods of the present invention for delivering the polypeptides of the invention. Liposomes are the preferred colloidal vehicle, and are composed of a phospholipid bilayer that may act as a carrier for both hydrophilic and hydrophobic medications. Liposomes can be made from, e.g., neutral lipids, charged phospholipids, and cholesterol. The addition of an amphophilic polymer such as polyethylene glycol (PEG) onto the surface of a liposome can slow the clearance of liposomes.

Administration of the polypeptides of the invention modified with PEG is also within the scope of the present invention. PEGs are polymers comprised of repeating ethylene oxide subunits with two terminal hydroxyl groups that can be chemically activated. PEG molecules come in a number of different configurations. PEG chains include linear and branched structures in which one or more PEG chains are joined with linkers such as lysine or triazine. PEGs may be

attached, preferably covalently, to a polypeptide of the invention at a single site or at multiple sites. Since branched-chain PEGs attach at single or fewer sites than do linear PEGs, branched PEGs may be less likely to interfere with the biologic activity of the native molecule than would the attachment of multiple small linear-chain PEGs, and so are preferred. Pharmaceutical formulations suitable for oral administration of proteins are described, e.g., in U.S. Patents 5,008,114; 5,505,962; 5,641,515; 5,681,811; 5,700,486; 5,766,633; 5,792,451; 5,853,748; 5,972,387; 5,976,569; and 6,051,561.

The polypeptides of the invention can also be delivered to an individual in need of treatment using gene therapy methods. Any of the methods for gene therapy available in the art can be used according to the present invention. Exemplary methods are described below.

In a preferred aspect, the therapeutic comprises a nucleic acid encoding a polypeptide of the invention that is part of an expression vector that expresses a polypeptide of the invention or a chimeric protein thereof in a suitable host. In particular, such a nucleic acid has a promoter operably linked to the polypeptide coding region, the promoter being inducible or constitutive, and, optionally, tissue-specific. In another particular embodiment, a nucleic acid molecule is used in which the polypeptide coding sequences and any other desired sequences are flanked by regions that promote homologous recombination at a desired site in the genome, thus providing for intrachromosomal expression of the desired nucleic acid.

Delivery of a nucleic acid encoding a polypeptide of the invention into a patient may be either direct, in which case the patient is directly exposed to the nucleic acid or nucleic acid-carrying vector, or indirect, in which case, cells are first transformed with the nucleic acid *in vitro*, then transplanted into the patient. These two approaches are known, respectively, as *in vivo* or *ex vivo* gene therapy.

In a specific embodiment, a nucleic acid encoding a polypeptide of the invention is directly administered *in vivo*, where it is expressed to produce the encoded product. This can be accomplished by any of numerous methods known in the art, e.g., by constructing it as part of an appropriate nucleic acid expression vector and administering it so that it becomes intracellular, e.g., by infection using a defective or attenuated retroviral or other viral vector (see, e.g., U.S. Pat. No. 4,980,286 and others mentioned *infra*), or by direct injection of naked DNA (see, e.g., Blezinger et al., *Nature Biotechnology*, 17, 343-348 (1999)) or by use of microparticle

bombardment (e.g., a gene gun; Biolistic, Dupont), or coating with lipids or cell-surface receptors or transfecting agents, encapsulation in liposomes (see, e.g., Chen et al., *Cancer Research*, 59, 3308-3312 (1999)), microparticles, or microcapsules, or by administering it in linkage to a peptide which is known to enter the nucleus, by administering it in linkage to a ligand subject to receptor-mediated endocytosis (see e.g., U.S. Patents 5,166,320; 5,728,399; 5,874,297; and 6,030,954) (which can be used to target cell types specifically expressing the receptors), etc.. In another embodiment, a nucleic acid-ligand complex can be formed in which the ligand comprises a fusogenic viral peptide to disrupt endosomes, allowing the nucleic acid to avoid lysosomal degradation. In yet another embodiment, the nucleic acid can be targeted *in vivo* for cell specific uptake and expression, by targeting a specific receptor (see, e.g., PCT Publications WO 92/06180; WO 92/22635; WO92/20316; WO93/14188; and WO 93/20221). Alternatively, the nucleic acid can be introduced intracellularly and incorporated within host cell DNA for expression, by homologous recombination (see, e.g., U.S. Patents 5,413,923; 5,416,260; and 5,574,205).

In a specific embodiment, a viral vector that contains a nucleic acid encoding a polypeptide of the invention is used. For example, a retroviral vector can be used (see, e.g., U.S. Patents 5,219,740; 5,604,090; and 5,834,182). These retroviral vectors have been modified to delete retroviral sequences that are not necessary for packaging of the viral genome and integration into host cell DNA. A nucleic acid encoding a polypeptide of the invention to be used in gene therapy is cloned into the vector, which facilitates delivery of the gene into a patient.

Adenoviruses are another type of viral vector that can be used in gene therapy. Adenovirus genomes are linear, double-stranded DNA molecules of approximately 36 kilobase pairs. Each extremity of the viral genome has a short sequence known as the inverted terminal repeat (or ITR), which is necessary for viral replication. The well-characterized molecular genetics of adenovirus render it an advantageous vector for gene transfer. Portions of the viral genome can be substituted with DNA of foreign origin. In addition, recombinant adenoviruses are structurally stable and no rearranged viruses are observed after extensive amplification.

Adenoviruses are especially attractive vehicles for delivering genes to respiratory epithelia. Adenoviruses naturally infect respiratory epithelia where they cause a mild disease. Other targets for adenovirus-based delivery systems are liver cells, the central nervous system, endothelial cells, and muscle. Adenoviruses have the advantage of being capable of infecting

non-dividing cells. Methods for conducting adenovirus-based gene therapy are described in e.g., U.S. Patents 5,824,544; 5,868,040; 5,871,722; 5,880,102; 5,882,877; 5,885,808; 5,932,210; 5,981,225; 5,994,106; 5,994,132; 5,994,134; 6,001,557; and 6,033,8843.

In a specific embodiment, the nucleic acid to be introduced for purposes of gene therapy using an adenoviral vector comprises an inducible promoter operably linked to the coding region, such that expression of the nucleic acid is controllable by controlling the presence or absence of the appropriate inducer of transcription.

The incorporation of genomic elements into the adenoviral vector may provide for enhanced expression of the DNA sequence encoding a polypeptide of the invention. Thus, in accordance with another aspect of the present invention, there is provided an adenoviral vector including at least one DNA sequence encoding a polypeptide of the invention, and at least one genomic element affecting the expression of such DNA sequence. The term "genomic element" is used as previously defined. Such genomic elements include, but are not limited to, introns, the 5' untranslated region, and the 3' untranslated region, and portions of the introns and 3' and 5' untranslated regions. The adenoviral vector may be as herein described. Promoters which control the DNA sequence may be selected from those described herein and from those known in the art.

The vector, consisting of infectious, but replication-defective, viral particles, which contain at least one DNA sequence encoding a polypeptide of the invention, is administered *in vivo* to a host in an amount effective to treat neovascularization or a tumour disease in the host. The host may be a mammalian host, including human and non-human primate hosts.

A viral vector can be administered in an amount of from about  $10^8$  plaque forming units (pfu) to about  $10^{14}$  plaque forming units, preferably from about  $10^8$  plaque forming units to about  $10^{11}$  plaque forming units, more preferably from about  $10^9$  plaque forming units to about  $10^{10}$  plaque forming units. Adenoviral vectors, in the quantities set out above, are preferred.

The infectious vector particles can be administered systemically, such as, for example, by intravenous administration (such as, for example, via peripheral vein injection) or administered via the portal vein, to the bile duct, intramuscularly, intraperitoneally, or intranasally. Alternatively,

infectious vector particles can be administered locally, by, e.g, intraocular injection or injection at the site of a tumour. Intraocular injection can be either into the anterior or posterior chamber of the eye, e.g., into the aqueous humor or vitreous humor. Alternatively, the injection can be subretinal, e.g., by injection of aliquots (e.g., 1 to 10 microliters per aliquot) of vector-containing solution behind the retina, after which the solution is absorbed and the infectious vector particles infect local cells of the ocular tissues and produce the active polypeptide. Such administration can comprise either a single injection, multiple injections administered on the same day, single injections administered over a period of weeks or months, or multiple injections administered over a period of weeks or months.

The vector particles may be administered in combination with a pharmaceutically acceptable carrier suitable for administration to a patient. The carrier may be a liquid carrier (for example, a saline solution), or a solid carrier, such as, for example, microcarrier beads.

Adeno-associated virus (AAV) has also been proposed for use in gene therapy, including e.g. endostatin gene therapy for tumours (see, e.g., Nguyen et al., Cancer Research, 58, 5673-5677 (1998)). Methods for producing and utilizing AAV are described, e.g., in U.S. Patents 5,173,414; 5,252,479; 5,552,311; 5,658,785; 5,763,416; 5,773,289; 5,843,742; 5,869,040; 5,942,496; and 5,948,675.

Other viral vectors which can be used in accordance with the present invention include especially gutless adenoviral vector, retroviral-based vectors such as Moloney murine leukemia virus and spleen necrosis virus, and other vectors derived from retroviruses such as Rous Sarcoma virus, Harvey sarcoma virus, avian leukosis virus, myeloproliferative sarcoma virus and mammary tumour virus. Lentiviral vectors are also preferred, especially bovine immunodeficiency viral vectors, such as those disclosed in PCT publication WO 01/44458.

Another approach to gene therapy involves transferring a gene to cells in tissue culture by such methods as electroporation, lipofection, calcium phosphate mediated transfection, or viral infection. Usually, the method of transfer includes the transfer of a selectable marker to the cells. The cells are then placed under selection to isolate those cells that have taken up and are expressing the transferred gene. Those cells are then delivered to a patient.

In this embodiment, the nucleic acid is introduced into a cell prior to administration *in vivo* of the resulting recombinant cell. Such introduction can be carried out by any method known in the art, including but not limited to transfection, electroporation, microinjection, infection with a viral or bacteriophage vector containing the nucleotide sequences, cell fusion, chromosome-mediated gene transfer, microcell-mediated gene transfer, spheroplast fusion, etc.. Numerous techniques are known in the art for the introduction of foreign genes into cells and may be used in accordance with the present invention, provided that the necessary developmental and physiological functions of the recipient cells are not disrupted. The technique should provide for the stable transfer of the nucleic acid to the cell, so that the nucleic acid is expressible by the cell and preferably heritable and expressible by its cell progeny.

Methods for the production and administration of such cells (i.e., those that produce high levels of protein from either endogenous or exogenous genes or nucleic acids) are described in, *inter alia*, U.S. Patent Nos. 5,641,670; 5,733,761; 5,968,502; 6,048,729; 6,054,288; 6,063,630; and 6,187,305. In a preferred embodiment, cells producing a polypeptide of the invention are delivered in microencapsulated form, e.g., in the form of cells microencapsulated in sodium alginate or calcium alginate poly L-lysine alginate (see, e.g., Read *et al.*, *Nature Biotechnology* 19, 29-34 (Jan 2001) and Joki *et al.*, *Nature Biotechnology* 19, 35-39 (Jan 2001)). The microencapsulated cells can be implanted proximally to the eye or the tumour, or at a site where a polypeptide of the invention produced by the cells will most quickly enter the subject's bloodstream, e.g., in the liver. The amount of cells envisioned for use depends on the desired effect, patient state, etc., and can be determined by one skilled in the art.

Cells into which a nucleic acid can be introduced for purposes of gene therapy encompass any desired, available cell type, and include but are not limited to epithelial cells, endothelial cells, keratinocytes, fibroblasts, muscle cells, hepatocytes; blood cells such as T lymphocytes, B lymphocytes, monocytes, macrophages, neutrophils, eosinophils, megakaryocytes, granulocytes; various stem or progenitor cells, in particular hematopoietic stem or progenitor cells, e.g., as obtained from bone marrow, umbilical cord blood, peripheral blood, fetal liver, etc..

In a preferred embodiment, the cell used for gene therapy is autologous to the patient.

In an embodiment in which recombinant cells are used in gene therapy, a nucleic acid encoding a polypeptide of the invention is introduced into the cells such that it is expressible by the cells or



their progeny, and the recombinant cells are then administered *in vivo* for therapeutic effect. In a specific embodiment, stem or progenitor cells are used. Any stem-and/or progenitor cells which can be isolated and maintained *in vitro* can potentially be used in accordance with this embodiment of the present invention. Such stem cells include but are not limited to hematopoietic stem cells (HSCs), stem cells of epithelial tissues such as the skin and the lining of the gut, embryonic heart muscle cells, liver stem cells (see, e.g., WO 94/08598), and neural stem cells.

Epithelial stem cells (ESCs) or keratinocytes can be obtained from tissues such as the skin and the lining of the gut by known procedures. In stratified epithelial tissue such as the skin, renewal occurs by mitosis of stem cells within the germinal layer, the layer closest to the basal lamina. Stem cells within the lining of the gut provide for a rapid renewal rate of this tissue. ESCs or keratinocytes obtained from the skin or lining of the gut of a patient or donor can be grown in tissue culture. If the ESCs are provided by a donor, a method for suppression of host versus graft reactivity (e.g., irradiation, drug or antibody administration to promote moderate immunosuppression) can also be used.

With respect to HSCs, any technique which provides for the isolation, propagation, and maintenance *in vitro* of HSCs can be used in this embodiment of the invention. Techniques by which this may be accomplished include (a) the isolation and establishment of HSC cultures from bone marrow cells isolated from the future host, or a donor, or (b) the use of previously established long-term HSC cultures, which may be allogeneic or xenogeneic. Non-autologous HSCs are used preferably in conjunction with a method of suppressing transplantation immune reactions of the future host/patient. In a particular embodiment of the present invention, human bone marrow cells can be obtained from the posterior iliac crest by needle aspiration (see, e.g., Kodo et al., 1984, J. Clin. Invest. 73:1377-1384). HSCs can be made highly enriched or in substantially pure form. This enrichment can be accomplished before, during, or after long-term culturing, and can be done by any techniques known in the art. Long-term cultures of bone marrow cells can be established and maintained by using, for example, modified Dexter cell culture techniques (Dexter et al., 1977, J. Cell Physiol. 91:335) or Witlock-Witte culture techniques (Witlock and Witte, 1982, Proc. Natl. Acad. Sci. USA 79:3608-3612).

It is understood, however, that the scope of the present invention is not to be limited to the specific embodiments described above. The invention may be practiced other than as particularly described and still be within the scope of the accompanying claims.

The pharmacological useful properties of the polypeptides of the present invention as therapeutics against tumour diseases and ocular neovascularization can be demonstrated for example by using known animal models for said diseases, such as those described e.g. in the Examples below, and applying the polypeptides, nucleic acid molecules or recombinant vectors of the present invention in these models.

Description of the Figures:

**Fig. 1:** Growth of A375 tumours in the flanks of immunosuppressive mice, measured as tumour weight 14 days after application of tumour cells.

*Control:* Tumours which develop within 14 days after application of  $1 \times 10^6$  A375 tumour cells, that are stably transfected with the empty pCDNA3 vector.

*s-EphB4:* Tumours which develop within 14 days after application of  $1 \times 10^6$  A375 tumour cells, which are stably transfected with the pCDNA3 vector containing the coding sequence for the extracellular domains of the Eph B4 receptor and which secrete the soluble s-EphB4 polypeptide.

*s-ephrinB2:* Tumours which develop within 14 days after application of  $1 \times 10^6$  A375 tumour cells, which are stably transfected with the pCDNA3 vector containing the coding sequence for the extracellular domains of the ephrin B2 ligand and which secrete the soluble s-ephrinB2 polypeptide.

**Fig. 2:** Growth of A375 tumours in the flanks of immunosuppressive mice.

Immunosuppressive mice are injected either in both flanks with A375 control tumour cells (animal group 1), or are injected in both flanks with A375 tumour cells which secrete s-EphB4 (animal group 2), or are injected in the left flank with A375 control tumour cells and in the right flank with A375 tumour cells which secrete s-EphB4 (animal group 3). The figure shows the tumour weight after 14 days (A) and the growth of the tumours (B). In the graph depicting the growth of tumours, for animal groups 1 and 2 the tumour volume of tumours from both flanks is determined.

Examples:

The following Examples serve to illustrate the invention without limiting its scope.

Example 1: Inhibition of tumour angiogenesis and tumour growth by expression of the soluble extracellular domains of murine Eph B4 (murine s-EphB4) or ephrin B2 (murine s-ephrinB2) in tumour cells.

Cloning of murine s-EphB4 and s-ephrinB2 eukaryotic expression vectors:

The nucleotide sequences coding for the soluble extracellular domains of murine Eph B4 (SEQ ID NO:1; nt 103-1719 of GeneBank Database Accession No. U06834) and murine ephrin B2 (SEQ ID NO:5; nt 161-847 of GeneBank Database Accession No. L38847) are amplified by means of sequence-specific oligonucleotide primers and Reverse Transcription Polymerase Chain Reaction (RT PCR) from isolated total RNA from mouse cerebral tissue or mouse kidney tissue, and modified with corresponding restriction cleavage sites as well as a stop codon. The amplified DNA is cloned by means of HindIII and BamHI into the vector pCDNA3 (Invitrogen) and the accuracy of the sequence is checked. The pCDNA3 vector contains the cytomegalovirus (CMV) promoter region and the G418 resistance gene. With this vector, the corresponding polypeptides s-EphB4 and s-ephrinB2 can be constitutively expressed in mammal cells under the control of the CMV promoter.

Generation of stably transfected tumour cells, which express either s-EphB4 (SEQ ID NO:2) or s-ephrinB2 (SEQ ID NO:6):

The recombinant vectors with the specific murine s-EphB4 and s-ephrinB2 nucleotide sequences, as well as the empty vector, are transfected with lipofectin (Gibco, LifeScience) into A375 human melanoma cells (ATCC No.: CRL-1619). Stable cell clones are generated by selection with 0.5 mg/ml G418 (Gibco, LifeScience). The expression of s-EphB4 and s-ephrinB2 is verified by Northern analysis and by a ligand receptor binding test [to this end, each 2 ml of conditioned tumour cell medium is concentrated by about five times (Amicon, Zentrifräp). 0.1 ml of concentrate is applied to a nitrocellulose membrane, and the membrane is subsequently incubated for one hour at room temperature in Tris-buffered saline (TBS) with 1% BSA. s-ephrinB2 is detected by incubation with biotinilated s-EphB4-Fc fusion protein (1 µg/ml; R&D Systems, Minneapolis, MN) and streptavidin coupled alkaline phosphatase. s-EphB4 is detected by incubation with biotinilated s-ephrinB2-Fc

fusion protein (1 µg/ml; R&D Systems, Minneapolis, MN) and streptavidin coupled alkaline phosphatase. Evidence of binding is found by enzymatic detection of the alkaline phosphatase]. In the proliferation assay, the transfected cell clones did not show differing growth kinetics (for the proliferation test, 8000 cells were seeded in a 24-well cell culture plate and the cell count was determined after 24, 48, 72 and 96 hours).

The influence of s-EphB4 and s-ephrinB2 on tumour growth and tumour angiogenesis:  $1 \times 10^6$  transfected A375 tumour cells are each injected intradermally into one or both flanks of immunosuppressive mice. Tumour growth is monitored over 14 days. After 14 days, the test animals are killed, the tumours prepared, weighed and examined histologically and by means of immunohistochemistry.

Monitoring of the body weight of the test animal, as well as observation of conspicuous behavioural patterns, are carried out to check for acute toxic side effects.

#### Results:

Animals which are injected with tumour cells (A375), which constitutively secrete either s-EphB4 or s-ephrinB2, develop considerably smaller tumours than animals which are injected with control cells, which are transfected with the empty vector (Fig. 1). The tumours, which are formed by A375 tumour cells and constitutively secrete either s-EphB4 or s-ephrinB2, have a disordered vessel structure compared to control tumours.

Animals that are injected with control tumour cells in both flanks develop tumours of equal size on both flanks. The kinetics of tumour growth is unchanged on both flanks (for the kinetics of tumour growth, the tumour volume (greatest diameter and perpendicular on this diameter) is measured by a slide gauge. The volume is determined by the factor  $0.5 \times \text{tumour diameter [mm]} \times \text{perpendicular}^2 [\text{mm}^2]$ ). Animals which are injected in both flanks with s-EphB4 secreting tumour cells, develop tumours of equal size on both sides, but the tumour size is significantly smaller than that of the corresponding control tumours. The kinetics of tumour growth is unchanged on both flanks.

Animals which are injected with control tumour cells in the left flank and s-EphB4 secreting tumour cells in the right flank, develop tumours of different size. On the left flank, considerably smaller control tumours develop, compared with the control tumours of animals which are injected with control cells in both flanks (Fig. 2 A, B).

In the groups of test animals, conspicuous behaviour patterns, such as motor disorders or hypermotor activities, are not observed, nor noticeable weight loss.

Example 2: Therapeutical application of recombinant s-EphB4 (SEQ ID NO:2) and s-ephrinB2 (SEQ ID NO:6).

The cDNA sequences coding for the soluble extracellular domains of the murine Eph B4 receptor (SEQ ID NO:1) or for those of the murine ephrin B2 ligand (SEQ ID NO:5), are cloned into a eukaryotic expression vector (for example pCDNA3, see also Example 1). "Chinese Hamster Ovarian" (CHO) cells are transfected with these recombinant vectors and stable cell clones are isolated by selection markers.

The expression levels of secreted soluble s-EphB4 protein and s-ephrinB2 protein from stably transfected CHO cell clones are determined by Western Blot Analysis (anti-EphB4 antibodies, Santa Cruz Ltd; anti-EphrinB2 antibodies, R&D Systems, Minneapolis, MN). Functionality is tested in a binding test, in which cell supernatants of stably transfected CHO cells are applied to a nitrocellulose membrane, the membrane is subsequently incubated for one hour at room temperature in TBS with 1% BSA. s-ephrinB2 is detected by incubation with biotinilated s-EphB4-Fc fusion protein (1 µg/ml, R&D Systems, Minneapolis, MN) and streptavidin coupled alkaline phosphatase. s-EphB4 is detected by incubation with biotinilated s-ephrinB2-Fc fusion protein (1 µg/ml, R&D Systems, Minneapolis, MN) and streptavidin coupled alkaline phosphatase. Evidence of binding is found by enzymatic detection of the alkaline phosphatase. CHO cell clones which noticeably secrete functionally active s-EphB4 protein or s-ephrinB2 protein are fermented and the recombinant proteins from the cell supernatant are purified. Purification is carried out by column chromatography with ConcavalinA affinity chromatography (Amersham/Pharmacia) followed by DEAE sepharose (Amersham/Pharmacia). Separation according to size is then carried out by gel filtration. The purified protein is dissolved in "Hanks buffered salt solution" (HBSS), sterile-filtered and stored in lyophilised state. Directly before therapeutical usage, the protein is dissolved in e.g. sterile water in a concentration of 2 mg/ml.

**Therapy test:**

A375 tumours are induced by intradermal injection of  $1 \times 10^6$  A375 melanoma cells into the flanks of immunosuppressive mice. Animals which develop a readily tangible tumour are treated by daily intraperitoneal injection of 0.1 mg/animal of recombinant protein and compared with tumour-bearing control animals which are given the same intraperitoneal

volume of HBSS. The success of therapy is determined by the tumour growth and histochemical and immunohistochemical characterization.

Monitoring of the body weight of the test animal, as well as observation of conspicuous behavioural patterns, are carried out to check for acute toxic side effects.

#### Results:

A375 tumour-bearing, immunosuppressive mice, which receive injections of recombinant, purified s-EphB4 polypeptid or s-ephrinB2 polypeptid, develop considerably smaller tumours with disorderd vessel structure compared to control animals which are given corresponding volumes of HBSS.

In the groups of test animals, conspicuous behaviour patterns, such as motor disorders or hypermotor activities, are not observed, nor noticeable weight loss.

#### Example 3: Generation and characterization of an adenoviral vector containing the DNA encoding for the soluble extracellular fragment of the human Eph B4 receptor.

Construction of adenoviral (Av3) vector expressing the extracellular domain of human Eph B4 receptor (human s-EphB4):

The cDNA encoding the extracellular domain of human Eph B4 is PCR-amplified from human fetus whole Marathon-Ready cDNA (Clontech Laboratories, Inc. Palo Alto, CA) with the primers; sense (5'- ATC ACT GCT AGC AAC TCA GTT CGG ATC CTA CC-3') and antisense (5'- ATC ATA GCT AGC TCA CTG CTC CCG CCA GCC CTC GC -3'). The PCR amplifies the region encoding the first 537 amino acids coding region of human Eph B4 (aa 1-537 of SEQ ID NO:4), including the signal peptide (the first 15 amino acids) and extracellular region of the mature Eph B4 receptor protein (16 to 537 amino acids). PCR is carried out with Pfu DNA polymerase (Stratagene, La Jolla, CA) in the presence of 1 µM primers as described, 0.4 mM dNTP, 1.5 mM MgCl<sub>2</sub> and 1 ng cDNA. The reaction is run for 35 cycles with the following conditions: 94 °C hot start for 1 min, 95 °C denaturation for 1 min, 62.5 to 67.5 °C with 0.5 °C thermal gradient annealing for 1 min, 72 °C extension for 3 min, and ended at 72 °C extension for 10 min. The 1674 base pairs (bp) s-EphB4 containing DNA fragments are pooled together, gel-purified and subcloned into pPCR-script plasmid according to the manufacturer's procedure (Stratagene, La Jolla, CA). The Nhe1 digested s-EphB4 DNA fragment is gel purified and cloned into the Spe1 site of the adenoviral shuttle plasmid, pAvsCvXlx [this vector contains the left end of the adenoviral vector

including the left inverted terminal repeat (ITR), a CMV promoter, an adenovirus major late promoter tripartite leader sequence, a multiple cloning site for inserting the transgene of interest, a SV40 early polyadenylation signal, and a loxP site], downstream of the simian CMV promoter and upstream of the SV40 polyadenylation signal to generate pAvCshs-EphB4-C16. The DNA sequence of the entire PCR-amplified region of the pAvCshs-EphB4-C16 is confirmed by sequence analysis by the Gene Therapy Core Technologies Molecular Core Laboratory.

The recombinant adenoviral vector Av3Cshs-EphB4 (with E1, E2a, and E3-deletions) is generated in the following manner. The pAvCshs-EphB4-C16 and pSQ3 (a 31'574 bp plasmid vector containing the right hand portion of the adenoviral vector genome and a loxP site that allows Cre-mediated recombination with pAvCshs-EphB4-C16) are first linearized with NotI and ClaI restriction enzymes, respectively. A transient transfection is performed on a 6-well plate containing  $5 \times 10^5$  S8 cells [S8 cells are derived from A549 cells (ATCC No. CCL-185) that have been transfected with plasmids containing the E1 and E2a regions, respectively, of the adenoviral vector (see also: Gorziglia et al., J. Virology 70, 4173-4178, 1996] per well using the LipofectAMINE PLUS reagent (Life Technologies, Rockville, MD) according to the manufacturer's procedure. The cells are cultured with media in the presence of 0.3  $\mu$ M of dexamethasone the day before transfection. The DNA-PLUS is prepared with 0.5  $\mu$ g of linearized pAvCshs-EphB4-C16, 1  $\mu$ g of linearized pSQ3 and 0.5  $\mu$ g of the expression plasmid pC-Cre3.1 encoding the Cre recombinase, in a 100- $\mu$ l Optimen (Life Technologies) with 6- $\mu$ l of PLUS reagent. After 15 min of incubation, the DNA-PLUS is mixed with 0.8-ml of 1 to 200-fold diluted lipofectAMINE Reagent. The DNA-PLUS-LipofectAMINE complex is added to the cells and incubated for 3 hours. The medium is then replenished with fresh medium. After seven to ten days incubation, cytopathic effect (CPE) is observed. The cells and the medium are harvested by scraping. A crude viral lysate (CVL) is prepared from 1 ml resuspended cell pellet by five cycles of freezing and thawing. The Av3Cshs-EphB4 vector is re-amplified in S8 cells with 0.3  $\mu$ M dexamethasone in Richter's CM medium containing 5% FBS until CPE was observed. The CVL is plaque-purified. The harvested single plaque free of replication competent adenovirus (RCA) is selected to generate purified vector.

**Av3Cshs-EphB4 research seed lot characterization:**

The research seedlot purified Av3Cshs-EphB4 vector is scaled up by the GTI Gene Therapy Core Technologies. The adenoviral vector titer (particles/ml) and biological titer (pfu/ml) are determined as described (Mittereder, N., March, K.L., and Trapnell, B.C. (1996) *J. Viro.* 70, 7498-7509) by GTI Gene Therapy Core Technologies. The ratio of total particles to infectious particles (particles/pfu) is calculated.

**Characterization of recombinant Av3Cshs-EphB4:**

Genomic DNA is prepared from the purified Av3Cshs-EphB4 vector using DNA isolation kit (Gentra Systems, Minneapolis, MN) according to the manufacturer's procedure. Various restriction digests by BamHI, Xba1, and XhoI of the purified viral DNA of Av3Cshs-EphB4, are compared on 1.0% agarose-TAE gel and stained with ethidium bromide to visualize the various digests for the verification of the correct genome structure and purity.

**Northern blot analysis:**

Hep3B (ATCC No. HB-8064) and A549 cells are transduced with Av3Cshs-EphB4, or the control Av3CsNull (same genome structure with no transgene in the expression cassette) vectors at the particles to cell ratio of 750. 72 hours after transduction, the cell pellets are harvested. Total RNA is isolated using the RNA-DNA STAT 60 (Tel-Test) extraction method. Northern blot analysis is carried out according the standard procedure. A 20 µg aliquot of total RNA is analyzed on 1% agarose formaldehyde/MOPS gel. RNA is transferred to a nylon membrane and prehybridized in the solution of 6.75% SDS, 0.5M NaPO<sub>4</sub>, pH 7.0, 1.0 mM EDTA, pH 8.0, 5 µg/ml BSA, and 100 µg/ml sheared, denatured salmon sperm DNA. The probe of 1368 bp s-EphB4 internal fragment is generated by EcoN1 digestion, gel purified and <sup>32</sup>P-labeled. The membrane is hybridized with a [<sup>32</sup>P]-labeled s-EphB4 internal probe at 68 °C overnight and washed in SSC/SDS containing buffers at 68 °C following the standard protocol. The membrane is exposed to an x-ray film at -70 °C overnight. The membrane is reprobed with [<sup>32</sup>P]-labeled Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) internal probe to verify equal amount of RNA loaded on each lane.

**SDS-PAGE and Western blot analysis:**

S8 and Hep3B cells are transduced with Av3Cshs-EphB4 or the control, Av3CsNull vector at the particles to cells ratio of 100 and 500, respectively. 72 hours post transduction, the



conditioned media are collected and subjected to SDS-PAGE analyses. The resolved protein bands are then electrotransferred to PVDF membrane using the Mini Trans-blot Electrophoretic Transfer Cell (Cat# 165-3165, Bio-Rad). Western blot analyses are carried out according to the following procedure. Briefly, the membrane blot is blocked with blocking buffer (5% powdered milk, 0.1% Tween-20 and PBS buffer) for 30 min, probed with 1:1000 diluted rabbit anti-EphB4 antiserum (generated at GTI through outside service by Affinity Bioreagents Inc., Golden, CO) in blocking buffer at 4 °C overnight, probed with 1:5000 diluted Horse Radish Peroxidase (HRP)-conjugated anti-rabbit IgG antibody in TPBS (0.1% Tween-20 and PBS buffer) (Amersham Pharmacia Biotech) at 4 °C for 1 hour and developed in chemiluminescence detection mixture (ECL kit, Amersham Pharmacia Biotech) for 1 min followed by x-ray film exposure.

Polyclonal anti-EphB4 antiserum is generated according to the following procedure. The rabbits are immunized with KLH conjugated synthetic peptide of (C)DGQWEELSLDEEQH (amino acids 39-43 of internal region of human Eph B4) followed by two boost injections every other month. Sera are collected and confirmed with reactivity directed against peptide antigen by ELISA and against Eph B4 protein by Western blot using purified recombinant Mouse EphB4/Fc chimera protein (Cat#446-B4, R&D Systems, Minneapolis, MN).

#### Solid phase EphB4-ephrinB2 binding assay:

The solid phase assay is carried out as described in the following procedure. Briefly, mouse EphB4/Fc chimera (Cat#446-B4-200, R&D Systems) is immobilized on a 96-well Nunc-Immuno™ plate with polySorp™ surface (Fisher) at 4 °C overnight. After 4 washes with TPBS (0.5% Tween-20 in DPBS), the plate is blocked with 10% FBS in DPBS for 30 min followed by 6 washes with TPBS. Various concentration of biotinylated mouse ephrinB2/Fc chimera (Cat#BT496, R&D Systems) in DPBS is then incubated with immobilized mouse EphB4/Fc chimera for 1 to 2 hours. After extensive washes with TPBS, the plate is incubated with 100 µl of 0.23 µg/ml of Avidin-HRP (Cat#21123, Pierce) for 1 hour followed by development in the Horseradish Peroxidase substrate mixture of 3, 3', 5, 5'-tetramethylbenzidine solution and hydrogen peroxide (Cat # 172-1064, Bio-Rad, Hercules, CA) for 10 to 30 min. The colored product is determined by absorbance at 415 nm by an ELISA reader. To determine the inhibitory effect of s-EphB4, 100 ng/ml (2.1 nM) of biotinylated mouse ephrinB2/Fc chimera is incubated for 30 min to 1 hour with increasing amounts of conditioned medium from untreated control S8 cells or S8 cells transduced with either null vector or s-EphB4 expressing vector prior to incubation of the mixture with immobilized

EphB4/Fc. Data is analyzed using Prism nonlinear regression curve fit to determine the best-fit curve and  $K_d$ , concentration of ligand required to reach half-maximal binding.

#### Matrigel Implant Assay:

Liquid Matrigel (a sterile extract of basement membrane proteins) is obtained from Fisher Scientific, Part number CB40234C, mixed with test substances and preloaded into a 1.0 ml tuberculin syringes with G27 needles. Male nude mice (Harlan) weighing about 20 g are anesthetized and 0.45 ml of the undiluted Matrigel alone or the undiluted Matrigel containing 1  $\mu\text{g/ml}$  of FGF-2 (Fibroblast Growth Factor 2) or 3  $\mu\text{g/ml}$  of VEGF (Vascular Endothelial Growth Factor; R&D Systems, Minneapolis, Minnesota) and  $5 \times 10^5$  S8 cells transduced at 100 MOI (multiplicity of infection) with either null vector or s-EphB4 expressing vector are injected subcutaneously into the dorsal caudal portion of the midline (1 injection/animal) using a G27 needle. For VEGF-induced angiogenesis, 20 U/ml heparin is also included. The Matrigel will rapidly form a solid gel that persists for over 7 days. On day 7, all animals are euthanized by  $\text{CO}_2$  and Matrigel plugs are harvested for histological analysis and hemoglobin measurement.

#### Hemoglobin Analysis:

Matrigel plugs are snap frozen with dry ice and dried over night. The dry weight is recorded and the plug rehydrated with 0.5 ml of 0.5% Tween-20 and water. After homogenization, the mixture is centrifuged at 14,000 rpm in a desktop centrifuge (Eppendorf 5415C, Brinkmann Instruments, Inc, Westbury, NY) for 30 min. Supernatant fluid is collected and its absorbance read at 405nm and converted to  $\mu\text{g}$  hemoglobin per mg Matrigel using a standard curve generated with hemoglobin standards. Data represent mean + SEM,  $n = 10$ . Experiments are done at least twice.

#### Results:

The gene therapy vector, Av3Cshs-EphB4 mediates s-EphB4 expression and secretion: Northern blotting analysis indicates that Av3Cshs-EphB4 transduced Hep3B and A549 cells express a 1.8 Kb mRNA transcript that specifically hybridizes to the Eph B4 cDNA probe. In Av3Cshs-EphB4 transduced S8 (without dexamethasone) and Hep3B cells, Western blot analysis also reveals a band with an apparent molecular mass of 85 kDa. This band is not observed in cells transduced with Av3CsNull or in untreated cells. The predicted molecular

mass of the expressed s-EphB4 protein is considerably smaller (57 kDa), suggesting that at least one of the three potential glycosylation sites is glycosylated.

**Inhibition of EphB4-ephrinB2 binding by s-EphB4:**

In the solid phase assay, dose-dependent saturation binding is demonstrated with the best curve-fit equation of  $\text{Absorbance} = \text{AbsMax} \times [\text{biotinylated ephrinB2}] / (\text{Kd} + [\text{biotinylated ephrinB2}])$  revealing a dissociation constant of  $1.16 \pm 0.24$  nM. AbsMax represents the absorbance at saturated biotinylated ephrinB2 concentration. The binding assay is then used to determine if s-EphB4 can effectively interfere with this binding. As shown in Table 1, conditioned media from untreated S8 cells and S8 transduced with null vector actually enhance the binding of Eph B4 with ephrin B2. In contrast, conditioned media prepared from Av3Cshs-EphB4 transduced S8 substantially inhibit the binding in a dose-dependent manner. These results indicate that this gene therapy vector can direct the expression of the extracellular domain of the Eph B4 receptor and can block the interaction of Eph B4 with its cognate partners.

**Adenoviral vector expressing the extracellular domain of Eph B4 can inhibit both VEGF and FGF-2 directed angiogenesis:**

As shown in Table 2, VEGF and FGF-2 are both able to substantially induce angiogenesis as measured by the increase in hemoglobin content in the Matrigel plug. However, FGF-2 is a more potent inducer of angiogenesis in this model (Table 2). Histological analysis reveals that the increase in hemoglobin content correlates with an increase in the number of blood vessels. The addition of Av3Cshs-EphB4 transduced cells substantially inhibits both VEGF- and FGF-2-induced angiogenesis. These results indicate that the soluble, extracellular domain of Eph B4 can inhibit angiogenesis induced by VEGF or FGF-2.

**Table 1: Inhibition of EphB4-ephrinB2 binding by gene therapy vector-mediated expression of the soluble, extracellular domain of Eph B4**

%Conditioned media	Absorbance at 415 nm		
	Untreated S8 cells	S8/Av3CsNull	S8/Av3Cshs-EphB4
0	0.63 ± 0.06	0.52 ± 0.11	0.60 ± 0.06
12.5	0.74 ± 0.06	0.55 ± 0.09	0.43 ± 0.07
25	0.89 ± 0.11	0.81 ± 0.14	0.41 ± 0.03
50	0.81 ± 0.15	0.76 ± 0.10	0.31 ± 0.02
100	0.97 ± 0.03	0.90 ± 0.26	0.29 ± 0.01

**Table 2: Inhibition of *in vivo* angiogenesis by gene therapy vector-mediated expression of the soluble, extracellular domain of Eph B4**

Treatment	Hemoglobin content/ Matrigel [µg/mg]	SEM	% Inhibition
Matrigel alone	0.48	0.06	
VEGF + null vector	4.24	0.46	
VEGF + s-EphB4	2.47	0.31	47%
FGF + null vector	12.85	1.68	
FGF + s-EphB4	9.01	1.26	31%

Example 4: Application of Av3Csms-EphB4 or Av3Csms-ephrinB2 vector in a mouse tumour model.

The recombinant adenoviral vectors Av3Csms-EphB4 and Av3CshsephrinB2 which contain the DNA encoding the soluble extracellular fragment of murine EphB4 and ephrinB2, respectively, are prepared analogously to the preparation of Av3Cshs-EphB4 as described in Example 3 above using the murine s-EphB4 and s-ephrinB4 cDNAs, respectively, instead of the human s-EphB4 cDNA.

**Gene therapy:**

A375 tumours are induced by intradermal injection of  $1 \times 10^6$  A375 melanoma cells into the flanks of immunosuppressive mice. Animals which develop a readily tangible tumour are treated by intravenous injection of Av3Csms-EphB4 or Av3CshsephrinB2 ( $1 \times 10^8$  "plaque forming units" (pfu)/animal –  $1 \times 10^9$  pfu/animal) and compared with tumour-bearing control animals which are treated with recombinant control adenoviruses (same genome structure with no transgene in the expression cassette) vectors. The success of therapy is determined by the tumour growth. The resulting tumours are further examined by histochemistry and immunohistochemistry.

Monitoring of the body weight of the test animal, as well as observation of conspicuous behavioural patterns, are carried out to check for acute toxic side effects.

**Results:**

A375 tumour-bearing, immunosuppressive mice which receive recombinant adenoviruses, which contain either the coding sequence for the extracellular domains of the murine Eph B4 receptor or that of the murine ephrin B2 ligand, develop substantially smaller tumours with a disordered vessel structure compared to control animals which receive control adenoviruses.

In the groups of test animals, conspicuous behaviour patterns, such as motor disorders or hypermotor activities, are not observed, nor noticeable weight loss.

What is claimed is:

1. An isolated soluble polypeptide which comprises an amino acid sequence of an extracellular region of an Eph B receptor or an essentially similar amino acid sequence thereof, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian ephrin B ligand.
2. A polypeptide according to claim 1 which comprises an amino acid sequence of an extracellular region of Eph B4 or an essentially similar amino acid sequence thereof, said polypeptide being capable of binding with high affinity to a mammalian ephrin B ligand.
3. A polypeptide according to claim 1 which comprises SEQ ID NO:2 or SEQ ID NO:4, or an essentially similar amino acid sequence thereof, said polypeptide being capable of binding with high affinity to a mammalian ephrin B ligand.
4. An isolated soluble polypeptide which comprises an amino acid sequence of an extracellular region of an ephrin B ligand or an essentially similar amino acid sequence thereof, said isolated soluble polypeptide being capable of binding with high affinity to a mammalian Eph B receptor.
5. A polypeptide according to claim 4 which comprises an amino acid sequence of an extracellular region of ephrin B2 or an essentially similar amino acid sequence thereof, said polypeptide being capable of binding with high affinity to a mammalian Eph B receptor.
6. A polypeptide according to claim 4 which comprises SEQ ID NO:6 or SEQ ID NO:8, or an essentially similar amino acid sequence thereof, said polypeptide being capable of binding with high affinity to a mammalian Eph B receptor.
7. A pharmaceutical composition comprising a polypeptide according to any one of claims 1 to 6.
8. Use of a polypeptide according to any one of claims 1 to 6 in the production of a pharmaceutical composition for the treatment of a tumour disease of mammals, including humans.

9. Use of a polypeptide according to any one of claims 1 to 6 in the production of a pharmaceutical composition for the treatment of ocular neovascularization in mammals, including humans.
10. An isolated nucleic acid molecule comprising a nucleotide sequence that encodes a polypeptide according to any one of claims 1 to 6.
11. A pharmaceutical composition comprising a nucleic acid molecule according to claim 10.
12. Use of a nucleic acid molecule according to claim 10 in the production of a pharmaceutical composition for the treatment of a tumour disease of mammals, including humans.
13. Use of a nucleic acid molecule according to claim 10 in the production of a pharmaceutical composition for the treatment of ocular neovascularization in mammals, including humans.
14. A recombinant vector comprising a nucleic acid molecule according to claim 10.
15. A recombinant vector according to claim 14, which is a recombinant viral vector.
16. A recombinant viral vector according to claim 15, which is a recombinant adenoviral vector.
17. A recombinant viral vector according to claim 15, which is a recombinant lentiviral vector.
18. A recombinant viral vector according to claim 15, which is a recombinant bovine immunodeficiency virus (BIV) vector.
19. A pharmaceutical composition comprising a recombinant vector according to any one of claims 14 to 18.

20. Use of a recombinant vector according to any one of claims 14 to 18 in the production of a pharmaceutical composition for the treatment of a disease of mammals, including humans, by means of gene therapy.


21. Use according to claim 20 for the treatment of a tumour disease of mammals, including humans.

22. Use according to claim 20 for the treatment of ocular neovascularization in mammals, including humans.

23. A host cell comprising a recombinant vector according to any one of claims 14 to 18.




1/2

**Fig. 1**

2/2

Fig. 2



- 1 -

## SEQUENCE LISTING

<110> Novartis AG  
 <120> Extracellular polypeptides of Eph B receptors and ephrin B ligands  
 and the corresponding nucleic acid molecules  
 <130> Case 4-31611A  
 <160> 8  
 <170> PatentIn version 3.0  
 <210> 1  
 <211> 1617  
 <212> DNA  
 <213> Mus musculus  
 <220>  
 <221> CDS  
 <222> (1)..(1617)  
 <400> 1  
 atg gag ctc cga gcg ctg ctg tgc tgg gct tcc ctc gcc act gct tta 48  
 Met Glu Leu Arg Ala Leu Leu Cys Trp Ala Ser Leu Ala Thr Ala Leu  
 1 5 10 15  
 gaa gag acc ctg ttg aac aca aaa ctg gaa acg gcg gat ctg aaa tgg 96  
 Glu Glu Thr Leu Leu Asn Thr Lys Leu Glu Thr Ala Asp Leu Lys Trp  
 20 25 30  
 gtg act tac cct cag gca gag ggc cag tgg gag gag cta agc ggc ctg 144  
 Val Thr Tyr Pro Gln Ala Glu Gly Gln Trp Glu Glu Leu Ser Gly Leu  
 35 40 45  
 gat gag gaa cag cac agc gtc cgc acc tat gag gtg tgc gac atg aag 192  
 Asp Glu Glu Gln His Ser Val Arg Thr Tyr Glu Val Cys Asp Met Lys  
 50 55 60  
 cgt cca ggg ggc cag gct cac tgg ctg cgc act ggc tgg gtc cca agg 240  
 Arg Pro Gly Gly Gln Ala His Trp Leu Arg Thr Gly Trp Val Pro Arg  
 65 70 75 80  
 cga ggt gct gtc cac gtg tat gcc acg ata cgc ttc acc atg atg gaa 288  
 Arg Gly Ala Val His Val Tyr Ala Thr Ile Arg Phe Thr Met Met Glu  
 85 90 95

- 2 -

tgc ctg tcc ctg ccg agg gcc agt cgc tcc tgc aag gag aca ttc act	336
Cys Leu Ser Leu Pro Arg Ala Ser Arg Ser Cys Lys Glu Thr Phe Thr	
100 105 110	
gtc ttc tat tac gag agc gaa cgt gat acg gcc acg gcc cat acg ccc	384
Val Phe Tyr Tyr Glu Ser Glu Arg Asp Thr Ala Thr Ala His Thr Pro	
115 120 125	
gcc tgg atg gag aac ccc tac atc aag gtg gac aca gtg gcc gca gaa	432
Ala Trp Met Glu Asn Pro Tyr Ile Lys Val Asp Thr Val Ala Ala Glu	
130 135 140	
cat ctg act cgg aag cgc cct gga gct gaa gcc aca ggg aaa gtt aat	480
His Leu Thr Arg Lys Arg Pro Gly Ala Glu Ala Thr Gly Lys Val Asn	
145 150 155 160	
atc aag acg ctg cgc ctg ggt cct ctc agc aaa gct ggc ttc tac ctg	528
Ile Lys Thr Leu Arg Leu Gly Pro Leu Ser Lys Ala Gly Phe Tyr Leu	
165 170 175	
gct ttc cag gac caa gga gcc tgc atg gct ctg ctc tcc ctg cat ctc	576
Ala Phe Gln Asp Gln Gly Ala Cys Met Ala Leu Leu Ser Leu His Leu	
180 185 190	
ttt tac aag aag tgc tcc tgg ctg atc acg aac ttg acc tac ttc ccc	624
Phe Tyr Lys Lys Cys Ser Trp Leu Ile Thr Asn Leu Thr Tyr Phe Pro	
195 200 205	
gag acg gtg cct cgg gag ctc gtg gtg ccg gtg gca ggt agc tgc gtg	672
Glu Thr Val Pro Arg Glu Leu Val Val Pro Val Ala Gly Ser Cys Val	
210 215 220	
gcc aac gcg gtc cct acc gcc aac ccc agc ccc agc ctc tac tgc cgg	720
Ala Asn Ala Val Pro Thr Ala Asn Pro Ser Pro Ser Leu Tyr Cys Arg	
225 230 235 240	
gaa gat ggt caa tgg gct gag cag cag gtc acg ggc tgc agc tgc gcg	768
Glu Asp Gly Gln Trp Ala Glu Gln Gln Val Thr Gly Cys Ser Cys Ala	
245 250 255	
cca ggg tac gag gct gcg gaa agc aac aaa gta tgc aga gcc tgt ggc	816
Pro Gly Tyr Glu Ala Ala Glu Ser Asn Lys Val Cys Arg Ala Cys Gly	
260 265 270	

- 3 -

cag gga acc ttc aag ccc caa ata gga gac gag tcc tgc ctg ccg tgc	864
Gln Gly Thr Phe Lys Pro Gln Ile Gly Asp Glu Ser Cys Leu Pro Cys	
275 280 285	
cca gcc aac agc cac tcg aat aac att ggg tct cct gtc tgc ctg tgt	912
Pro Ala Asn Ser His Ser Asn Asn Ile Gly Ser Pro Val Cys Leu Cys	
290 295 300	
cga att ggg tat tac cgg gcc cgc tca gac ccc cgg agt tca cct tgc	960
Arg Ile Gly Tyr Tyr Arg Ala Arg Ser Asp Pro Arg Ser Ser Pro Cys	
305 310 315 320	
act acc cca ccc tct gct cca aga agc gtg gtt cac cat ttg aat ggt	1008
Thr Thr Pro Pro Ser Ala Pro Arg Ser Val Val His His Leu Asn Gly	
325 330 335	
tcc acc ctg cgc ctg gag tgg agt gct ccc ctt gag tcc gga agg ccg	1056
Ser Thr Leu Arg Leu Glu Trp Ser Ala Pro Leu Glu Ser Gly Arg Pro	
340 345 350	
aga gac ctc act tat gct gtt cgc tgc cga gag tgc cgt cct ggg ggt	1104
Arg Asp Leu Thr Tyr Ala Val Arg Cys Arg Glu Cys Arg Pro Gly Gly	
355 360 365	
tcc tgc ttg ccc tgt ggg ggc gac atg acc ttc gac ccc ggt cct cga	1152
Ser Cys Leu Pro Cys Gly Gly Asp Met Thr Phe Asp Pro Gly Pro Arg	
370 375 380	
gac ctg gtt gag cgc tgg gtg gca atc cga ggg ctg cgt cct gat gtc	1200
Asp Leu Val Glu Arg Trp Val Ala Ile Arg Gly Leu Arg Pro Asp Val	
385 390 395 400	
acc tat acc ttt gag gtt gct gct ttg aat ggt gtg tct acc tta gcc	1248
Thr Tyr Thr Phe Glu Val Ala Ala Leu Asn Gly Val Ser Thr Leu Ala	
405 410 415	
act gga cca cct cct ttt gag cct gtc aat gtc acc act gac cgt gag	1296
Thr Gly Pro Pro Pro Phe Glu Pro Val Asn Val Thr Thr Asp Arg Glu	
420 425 430	
gtg cct cct gca gtg tct gac atc cga gtg act cgg tcg tca ccc agc	1344
Val Pro Pro Ala Val Ser Asp Ile Arg Val Thr Arg Ser Ser Pro Ser	
435 440 445	

- 4 -

agc ttg atc ctg tca tgg gct atc ccc aga gca ccc agt ggg gcc gtg 1392  
 Ser Leu Ile Leu Ser Trp Ala Ile Pro Arg Ala Pro Ser Gly Ala Val

450 455 460

ctg gac tac gag gtc aag tat cat gag aag ggc gca gag ggc ccc agc 1440

Leu Asp Tyr Glu Val Lys Tyr His Glu Lys Gly Ala Glu Gly Pro Ser

465 470 475 480

agt gtt cgt ttc ctg aag aca tca gaa aac cga gct gag ctc cgg ggg 1488

Ser Val Arg Phe Leu Lys Thr Ser Glu Asn Arg Ala Glu Leu Arg Gly

485 490 495

ctg aag cgg gga gcc agc tat ctg gtc cag gta cgc gca cgg tcc gag 1536

Leu Lys Arg Gly Ala Ser Tyr Leu Val Gln Val Arg Ala Arg Ser Glu

500 505 510

gct ggc tac ggt ccc ttc ggc cag gag cat cac agt cag act caa ctg 1584

Ala Gly Tyr Gly Pro Phe Gly Gln Glu His His Ser Gln Thr Gln Leu

515 520 525

gat gag agc gag agc tgg cgg gag cag ctg gcc 1617

Asp Glu Ser Glu Ser Trp Arg Glu Gln Leu Ala

530 535

<210> 2

<211> 539

<212> PRT

<213> Mus musculus

<400> 2

Met Glu Leu Arg Ala Leu Leu Cys Trp Ala Ser Leu Ala Thr Ala Leu

1 5 10 15

Glu Glu Thr Leu Leu Asn Thr Lys Leu Glu Thr Ala Asp Leu Lys Trp

20 25 30

Val Thr Tyr Pro Gln Ala Glu Gly Gln Trp Glu Glu Leu Ser Gly Leu

35 40 45

Asp Glu Glu Gln His Ser Val Arg Thr Tyr Glu Val Cys Asp Met Lys

50 55 60

- 5 -

Arg	Pro	Gly	Gly	Gln	Ala	His	Trp	Leu	Arg	Thr	Gly	Trp	Val	Pro	Arg	65	70	75	80
Arg	Gly	Ala	Val	His	Val	Tyr	Ala	Thr	Ile	Arg	Phe	Thr	Met	Met	Glu	85	90	95	
Cys	Leu	Ser	Leu	Pro	Arg	Ala	Ser	Arg	Ser	Cys	Lys	Glu	Thr	Phe	Thr	100	105	110	
Val	Phe	Tyr	Tyr	Glu	Ser	Glu	Arg	Asp	Thr	Ala	Thr	Ala	His	Thr	Pro	115	120	125	
Ala	Trp	Met	Glu	Asn	Pro	Tyr	Ile	Lys	Val	Asp	Thr	Val	Ala	Ala	Glu	130	135	140	
His	Leu	Thr	Arg	Lys	Arg	Pro	Gly	Ala	Glu	Ala	Thr	Gly	Lys	Val	Asn	145	150	155	160
Ile	Lys	Thr	Leu	Arg	Leu	Gly	Pro	Leu	Ser	Lys	Ala	Gly	Phe	Tyr	Leu	165	170	175	
Ala	Phe	Gln	Asp	Gln	Gly	Ala	Cys	Met	Ala	Leu	Leu	Ser	Leu	His	Leu	180	185	190	
Phe	Tyr	Lys	Lys	Cys	Ser	Trp	Leu	Ile	Thr	Asn	Leu	Thr	Tyr	Phe	Pro	195	200	205	
Glu	Thr	Val	Pro	Arg	Glu	Leu	Val	Val	Pro	Val	Ala	Gly	Ser	Cys	Val	210	215	220	
Ala	Asn	Ala	Val	Pro	Thr	Ala	Asn	Pro	Ser	Pro	Ser	Leu	Tyr	Cys	Arg	225	230	235	240
Glu	Asp	Gly	Gln	Trp	Ala	Glu	Gln	Gln	Val	Thr	Gly	Cys	Ser	Cys	Ala	245	250	255	
Pro	Gly	Tyr	Glu	Ala	Ala	Glu	Ser	Asn	Lys	Val	Cys	Arg	Ala	Cys	Gly	260	265	270	
Gln	Gly	Thr	Phe	Lys	Pro	Gln	Ile	Gly	Asp	Glu	Ser	Cys	Leu	Pro	Cys	275	280	285	

- 6 -

Pro Ala Asn Ser His Ser Asn Asn Ile Gly Ser Pro Val Cys Leu Cys  
 290 295 300

Arg Ile Gly Tyr Tyr Arg Ala Arg Ser Asp Pro Arg Ser Ser Pro Cys  
 305 310 315 320

Thr Thr Pro Pro Ser Ala Pro Arg Ser Val Val His His Leu Asn Gly  
 325 330 335

Ser Thr Leu Arg Leu Glu Trp Ser Ala Pro Leu Glu Ser Gly Arg Pro  
 340 345 350

Arg Asp Leu Thr Tyr Ala Val Arg Cys Arg Glu Cys Arg Pro Gly Gly  
 355 360 365

Ser Cys Leu Pro Cys Gly Gly Asp Met Thr Phe Asp Pro Gly Pro Arg  
 370 375 380

Asp Leu Val Glu Arg Trp Val Ala Ile Arg Gly Leu Arg Pro Asp Val  
 385 390 395 400

Thr Tyr Thr Phe Glu Val Ala Ala Leu Asn Gly Val Ser Thr Leu Ala  
 405 410 415

Thr Gly Pro Pro Pro Phe Glu Pro Val Asn Val Thr Thr Asp Arg Glu  
 420 425 430

Val Pro Pro Ala Val Ser Asp Ile Arg Val Thr Arg Ser Ser Pro Ser  
 435 440 445

Ser Leu Ile Leu Ser Trp Ala Ile Pro Arg Ala Pro Ser Gly Ala Val  
 450 455 460

Leu Asp Tyr Glu Val Lys Tyr His Glu Lys Gly Ala Glu Gly Pro Ser  
 465 470 475 480

Ser Val Arg Phe Leu Lys Thr Ser Glu Asn Arg Ala Glu Leu Arg Gly  
 485 490 495

Leu Lys Arg Gly Ala Ser Tyr Leu Val Gln Val Arg Ala Arg Ser Glu  
 500 505 510



- 7 -

Ala Gly Tyr Gly Pro Phe Gly Gln Glu His His Ser Gln Thr Gln Leu  
 515 520 525

Asp Glu Ser Glu Ser Trp Arg Glu Gln Leu Ala  
 530 535

<210> 3  
 <211> 1617  
 <212> DNA  
 <213> Homo sapiens  
 <220>  
 <221> CDS  
 <222> (1)..(1617)

<400> 3  
 atg gag ctc cgg gtg ctg ctc tgc tgg gct tgc ttg gcc gca gct ttg 48  
 Met Glu Leu Arg Val Leu Leu Cys Trp Ala Ser Leu Ala Ala Ala Leu  
 1 5 10 15  
 gaa gag acc ctg ctg aac aca aaa ttg gaa act gct gat ctg aag tgg 96  
 Glu Glu Thr Leu Leu Asn Thr Lys Leu Glu Thr Ala Asp Leu Lys Trp  
 20 25 30  
 gtg aca ttc cct cag gtg gac ggg cag tgg gag gaa ctg agc ggc ctg 144  
 Val Thr Phe Pro Gln Val Asp Gly Gln Trp Glu Glu Leu Ser Gly Leu  
 35 40 45  
 gat gag gaa cag cac agc gtg cgc acc tac gaa gtg tgt gaa gtg cag 192  
 Asp Glu Glu Gln His Ser Val Arg Thr Tyr Glu Val Cys Glu Val Gln  
 50 55 60  
 cgt gcc ccg ggc cag gcc cac tgg ctt cgc aca ggt tgg gtc cca cgg 240  
 Arg Ala Pro Gly Gln Ala His Trp Leu Arg Thr Gly Trp Val Pro Arg  
 65 70 75 80  
 cgg ggc gcc gtc cac gtg tac gcc acg ctg cgc ttc acc atg ctc gag 288  
 Arg Gly Ala Val His Val Tyr Ala Thr Leu Arg Phe Thr Met Leu Glu  
 85 90 95  
 tgc ctg tcc ctg cct cgg gct ggg cgc tcc tgc aag gag acc ttc acc 336  
 Cys Leu Ser Leu Pro Arg Ala Gly Arg Ser Cys Lys Glu Thr Phe Thr  
 100 105 110

- 8 -

gtc ttc tac tat gag agc gat gcg gac acg gcc acg gcc ctc acg cca	384
Val Phe Tyr Tyr Glu Ser Asp Ala Asp Thr Ala Thr Ala Leu Thr Pro	
115 120 125	
gcc tgg atg gag aac ccc tac atc aag gtg gac acg gtg gcc gcg gag	432
Ala Trp Met Glu Asn Pro Tyr Ile Lys Val Asp Thr Val Ala Ala Glu	
130 135 140	
cat ctc acc cgg aag cgc cct ggg gcc gag gcc acc ggg aag gtg aat	480
His Leu Thr Arg Lys Arg Pro Gly Ala Glu Ala Thr Gly Lys Val Asn	
145 150 155 160	
gtc aag acg ctg cgt ctg gga ccg ctc agc aag gct ggc ttc tac ctg	528
Val Lys Thr Leu Arg Leu Gly Pro Leu Ser Lys Ala Gly Phe Tyr Leu	
165 170 175	
gcc ttc cag gac cag ggt gcc tgc atg gcc ctg cta tcc ctg cac ctc	576
Ala Phe Gln Asp Gln Gly Ala Cys Met Ala Leu Leu Ser Leu His Leu	
180 185 190	
ttc tac aaa aag tgc gcc cag ctg act gtg aac ctg act cga ttc ccg	624
Phe Tyr Lys Lys Cys Ala Gln Leu Thr Val Asn Leu Thr Arg Phe Pro	
195 200 205	
gag act gtg cct cgg gag ctg gtt gtg ccc gtg gcc ggt agc tgc gtg	672
Glu Thr Val Pro Arg Glu Leu Val Val Pro Val Ala Gly Ser Cys Val	
210 215 220	
gtg gat gcc gtc ccc gcc cct ggc ccc agc ccc agc ctc tac tgc cgt	720
Val Asp Ala Val Pro Ala Pro Gly Pro Ser Pro Ser Leu Tyr Cys Arg	
225 230 235 240	
gag gat ggc cag tgg gcc gaa cag ccg gtc acg ggc tgc agc tgt gct	768
Glu Asp Gly Gln Trp Ala Glu Gln Pro Val Thr Gly Cys Ser Cys Ala	
245 250 255	
ccg ggg ttc gag gca gct gag ggg aac acc aag tgc cga gcc tgt gcc	816
Pro Gly Phe Glu Ala Ala Glu Gly Asn Thr Lys Cys Arg Ala Cys Ala	
260 265 270	
cag ggc acc ttc aag ccc ctg tca gga gaa ggg tcc tgc cag cca tgc	864
Gln Gly Thr Phe Lys Pro Leu Ser Gly Glu Gly Ser Cys Gln Pro Cys	
275 280 285	

- 9 -

cca gcc aat agc cac tct aac acc att gga tct gcc gtc tgc cag tgc	912
Pro Ala Asn Ser His Ser Asn Thr Ile Gly Ser Ala Val Cys Gln Cys	
290 295 300	
cgc gtc ggg gac ttc cgg gca cgc aca gac ccc cgg ggt gca ccc tgc	960
Arg Val Gly Asp Phe Arg Ala Arg Thr Asp Pro Arg Gly Ala Pro Cys	
305 310 315 320	
acc acc cct cct tgc gct ccg cgg agc gtg gtt tcc cgc ctg aac ggc	1008
Thr Thr Pro Pro Ser Ala Pro Arg Ser Val Val Ser Arg Leu Asn Gly	
325 330 335	
tcc tcc ctg cac ctg gaa tgg agt gcc ccc ctg gag tct ggt ggc cga	1056
Ser Ser Leu His Leu Glu Trp Ser Ala Pro Leu Glu Ser Gly Gly Arg	
340 345 350	
gag gac ctc acc tac gcc ctc cgc tgc cgg gag tgc cga ccc gga ggc	1104
Glu Asp Leu Thr Tyr Ala Leu Arg Cys Arg Glu Cys Arg Pro Gly Gly	
355 360 365	
tcc tgt gcg ccc tgc ggg gga gac ctg act ttt gac ccc ggc ccc cgg	1152
Ser Cys Ala Pro Cys Gly Gly Asp Leu Thr Phe Asp Pro Gly Pro Arg	
370 375 380	
gac ctg gtg gag ccc tgg gtg gtg gtt cga ggg cta cgt ccg gac ttc	1200
Asp Leu Val Glu Pro Trp Val Val Val Arg Gly Leu Arg Pro Asp Phe	
385 390 395 400	
acc tat acc ttt gag gtc act gca ttg aac ggg gta tcc tcc tta gcc	1248
Thr Tyr Thr Phe Glu Val Thr Ala Leu Asn Gly Val Ser Ser Leu Ala	
405 410 415	
acg ggg ccc gtc cca ttt gag cct gtc aat gtc acc act gac cga gag	1296
Thr Gly Pro Val Pro Phe Glu Pro Val Asn Val Thr Thr Asp Arg Glu	
420 425 430	
gta cct cct gca gtg tct gac atc cgg gtg acg cgg tcc tca ccc agc	1344
Val Pro Pro Ala Val Ser Asp Ile Arg Val Thr Arg Ser Ser Pro Ser	
435 440 445	
agc ttg agc ctg gcc tgg gct gtt ccc cgg gca ccc agt ggg gcg tgg	1392
Ser Leu Ser Leu Ala Trp Ala Val Pro Arg Ala Pro Ser Gly Ala Trp	
450 455 460	

- 10 -

ctg gac tac gag gtc aaa tac cat gag aag ggc gcc gag ggt ccc agc 1440  
 Leu Asp Tyr Glu Val Lys Tyr His Glu Lys Gly Ala Glu Gly Pro Ser  
 465 470 475 480

agc gtg cgg ttc ctg aag acg tca gaa aac cgg gca gag ctg cgg ggg 1488  
 Ser Val Arg Phe Leu Lys Thr Ser Glu Asn Arg Ala Glu Leu Arg Gly  
 485 490 495

ctg aag cgg gga gcc agc tac ctg gtg cag gta cgg gcg cgc tct gag 1536  
 Leu Lys Arg Gly Ala Ser Tyr Leu Val Gln Val Arg Ala Arg Ser Glu  
 500 505 510

gcc ggc tac ggg ccc ttc ggc cag gaa cat cac agc cag acc caa ctg 1584  
 Ala Gly Tyr Gly Pro Phe Gly Gln Glu His His Ser Gln Thr Gln Leu  
 515 520 525

gat gag agc gag ggc tgg cgg gag cag ctg gcc 1617  
 Asp Glu Ser Glu Gly Trp Arg Glu Gln Leu Ala  
 530 535

<210> 4

<211> 539

<212> PRT

<213> Homo sapiens

<400> 4

Met Glu Leu Arg Val Leu Leu Cys Trp Ala Ser Leu Ala Ala Ala Leu  
 1 5 10 15

Glu Glu Thr Leu Leu Asn Thr Lys Leu Glu Thr Ala Asp Leu Lys Trp  
 20 25 30

Val Thr Phe Pro Gln Val Asp Gly Gln Trp Glu Glu Leu Ser Gly Leu  
 35 40 45

Asp Glu Glu Gln His Ser Val Arg Thr Tyr Glu Val Cys Glu Val Gln  
 50 55 60

Arg Ala Pro Gly Gln Ala His Trp Leu Arg Thr Gly Trp Val Pro Arg  
 65 70 75 80

- 11 -

Arg Gly Ala Val His Val Tyr Ala Thr Leu Arg Phe Thr Met Leu Glu			
85	90	95	
Cys Leu Ser Leu Pro Arg Ala Gly Arg Ser Cys Lys Glu Thr Phe Thr			
100	105	110	
Val Phe Tyr Tyr Glu Ser Asp Ala Asp Thr Ala Thr Ala Leu Thr Pro			
115	120	125	
Ala Trp Met Glu Asn Pro Tyr Ile Lys Val Asp Thr Val Ala Ala Glu			
130	135	140	
His Leu Thr Arg Lys Arg Pro Gly Ala Glu Ala Thr Gly Lys Val Asn			
145	150	155	160
Val Lys Thr Leu Arg Leu Gly Pro Leu Ser Lys Ala Gly Phe Tyr Leu			
165	170	175	
Ala Phe Gln Asp Gln Gly Ala Cys Met Ala Leu Leu Ser Leu His Leu			
180	185	190	
Phe Tyr Lys Lys Cys Ala Gln Leu Thr Val Asn Leu Thr Arg Phe Pro			
195	200	205	
Glu Thr Val Pro Arg Glu Leu Val Val Pro Val Ala Gly Ser Cys Val			
210	215	220	
Val Asp Ala Val Pro Ala Pro Gly Pro Ser Pro Ser Leu Tyr Cys Arg			
225	230	235	240
Glu Asp Gly Gln Trp Ala Glu Gln Pro Val Thr Gly Cys Ser Cys Ala			
245	250	255	
Pro Gly Phe Glu Ala Ala Glu Gly Asn Thr Lys Cys Arg Ala Cys Ala			
260	265	270	
Gln Gly Thr Phe Lys Pro Leu Ser Gly Glu Gly Ser Cys Gln Pro Cys			
275	280	285	
Pro Ala Asn Ser His Ser Asn Thr Ile Gly Ser Ala Val Cys Gln Cys			
290	295	300	

- 12 -

Arg Val Gly Asp Phe Arg Ala Arg Thr Asp Pro Arg Gly Ala Pro Cys  
 305 310 315 320

Thr Thr Pro Pro Ser Ala Pro Arg Ser Val Val Ser Arg Leu Asn Gly  
 325 330 335

Ser Ser Leu His Leu Glu Trp Ser Ala Pro Leu Glu Ser Gly Gly Arg  
 340 345 350

Glu Asp Leu Thr Tyr Ala Leu Arg Cys Arg Glu Cys Arg Pro Gly Gly  
 355 360 365

Ser Cys Ala Pro Cys Gly Gly Asp Leu Thr Phe Asp Pro Gly Pro Arg  
 370 375 380

Asp Leu Val Glu Pro Trp Val Val Val Arg Gly Leu Arg Pro Asp Phe  
 385 390 395 400

Thr Tyr Thr Phe Glu Val Thr Ala Leu Asn Gly Val Ser Ser Leu Ala  
 405 410 415

Thr Gly Pro Val Pro Phe Glu Pro Val Asn Val Thr Thr Asp Arg Glu  
 420 425 430

Val Pro Pro Ala Val Ser Asp Ile Arg Val Thr Arg Ser Ser Pro Ser  
 435 440 445

Ser Leu Ser Leu Ala Trp Ala Val Pro Arg Ala Pro Ser Gly Ala Trp  
 450 455 460

Leu Asp Tyr Glu Val Lys Tyr His Glu Lys Gly Ala Glu Gly Pro Ser  
 465 470 475 480

Ser Val Arg Phe Leu Lys Thr Ser Glu Asn Arg Ala Glu Leu Arg Gly  
 485 490 495

Leu Lys Arg Gly Ala Ser Tyr Leu Val Gln Val Arg Ala Arg Ser Glu  
 500 505 510

Ala Gly Tyr Gly Pro Phe Gly Gln Glu His His Ser Gln Thr Gln Leu  
 515 520 525

- 13 -

Asp Glu Ser Glu Gly Trp Arg Glu Gln Leu Ala  
 530 535

<210> 5  
 <211> 687  
 <212> DNA  
 <213> Mus musculus  
 <220>  
 <221> CDS  
 <222> (1)..(687)

<400> 5  
 atg gcc atg gcc cgg tcc agg agg gac tct gtg tgg aag tac tgt tgg 48  
 Met Ala Met Ala Arg Ser Arg Arg Asp Ser Val Trp Lys Tyr Cys Trp  
 1 5 10 15  
 gga ctt ttg atg gtt ttg tgc aga act gcg atc tcc aga tcg ata gtt 96  
 Gly Leu Leu Met Val Leu Cys Arg Thr Ala Ile Ser Arg Ser Ile Val  
 20 25 30  
 tta gag cct atc tac tgg aat tcc tcg aac tcc aaa ttt cta ccc gga 144  
 Leu Glu Pro Ile Tyr Trp Asn Ser Ser Asn Ser Lys Phe Leu Pro Gly  
 35 40 45  
 caa ggc ctg gta cta tac cca cag ata gga gac aaa ttg gat att att 192  
 Gln Gly Leu Val Leu Tyr Pro Gln Ile Gly Asp Lys Leu Asp Ile Ile  
 50 55 60  
 tgc ccc aaa gtg gac tct aaa act gtt ggc cag tat gaa tat tat aaa 240  
 Cys Pro Lys Val Asp Ser Lys Thr Val Gly Gln Tyr Glu Tyr Tyr Lys  
 65 70 75 80  
 gtt tat atg gtt gat aaa gac caa gca gac aga tgc aca att aag aag 288  
 Val Tyr Met Val Asp Lys Asp Gln Ala Asp Arg Cys Thr Ile Lys Lys  
 85 90 95  
 gag aat acc ccg ctg ctc aac tgt gcc aga cca gac caa gat gtg aaa 336  
 Glu Asn Thr Pro Leu Leu Asn Cys Ala Arg Pro Asp Gln Asp Val Lys  
 100 105 110

- 14 -

ttc acc atc aag ttt caa gaa ttc agc cct aac ctc tgg ggt cta gaa 384  
 Phe Thr Ile Lys Phe Gln Glu Phe Ser Pro Asn Leu Trp Gly Leu Glu  
 115 120 125

ttt cag aag aac aaa gat tac tac att ata tct aca tca aat ggg tct 432  
 Phe Gln Lys Asn Lys Asp Tyr Tyr Ile Ile Ser Thr Ser Asn Gly Ser  
 130 135 140

ttg gag ggc ctg gat aac cag gag gga ggg gtg tgc cag aca aga gcc 480  
 Leu Glu Gly Leu Asp Asn Gln Glu Gly Gly Val Cys Gln Thr Arg Ala  
 145 150 155 160

atg aag atc ctc atg aaa gtt gga caa gat gca agt tct gct gga tca 528  
 Met Lys Ile Leu Met Lys Val Gly Gln Asp Ala Ser Ser Ala Gly Ser  
 165 170 175

gcc agg aat cac ggt cca aca aga cgt cca gag cta gaa gct ggt aca 576  
 Ala Arg Asn His Gly Pro Thr Arg Arg Pro Glu Leu Glu Ala Gly Thr  
 180 185 190

aat ggg aga agt tca aca aca agt ccc ttt gtg aag cca aat cca ggt 624  
 Asn Gly Arg Ser Ser Thr Thr Ser Pro Phe Val Lys Pro Asn Pro Gly  
 195 200 205

tct agc acc gat ggc aac agc gcg ggg cat tcc ggg aac aat ctc ctg 672  
 Ser Ser Thr Asp Gly Asn Ser Ala Gly His Ser Gly Asn Asn Leu Leu  
 210 215 220

ggt tcc gaa gtg gcc 687  
 Gly Ser Glu Val Ala  
 225

<210> 6  
 <211> 229  
 <212> PRT  
 <213> Mus musculus

<400> 6  
 Met Ala Met Ala Arg Ser Arg Arg Asp Ser Val Trp Lys Tyr Cys Trp  
 1 5 10 15



- 15 -

Gly Leu Leu Met Val Leu Cys Arg Thr Ala Ile Ser Arg Ser Ile Val  
 20 25 30

Leu Glu Pro Ile Tyr Trp Asn Ser Ser Asn Ser Lys Phe Leu Pro Gly  
 35 40 45

Gln Gly Leu Val Leu Tyr Pro Gln Ile Gly Asp Lys Leu Asp Ile Ile  
 50 55 60

Cys Pro Lys Val Asp Ser Lys Thr Val Gly Gln Tyr Glu Tyr Tyr Lys  
 65 70 75 80

Val Tyr Met Val Asp Lys Asp Gln Ala Asp Arg Cys Thr Ile Lys Lys  
 85 90 95

Glu Asn Thr Pro Leu Leu Asn Cys Ala Arg Pro Asp Gln Asp Val Lys  
 100 105 110

Phe Thr Ile Lys Phe Gln Glu Phe Ser Pro Asn Leu Trp Gly Leu Glu  
 115 120 125

Phe Gln Lys Asn Lys Asp Tyr Tyr Ile Ile Ser Thr Ser Asn Gly Ser  
 130 135 140

Leu Glu Gly Leu Asp Asn Gln Glu Gly Gly Val Cys Gln Thr Arg Ala  
 145 150 155 160

Met Lys Ile Leu Met Lys Val Gly Gln Asp Ala Ser Ser Ala Gly Ser  
 165 170 175

Ala Arg Asn His Gly Pro Thr Arg Arg Pro Glu Leu Glu Ala Gly Thr  
 180 185 190

Asn Gly Arg Ser Ser Thr Thr Ser Pro Phe Val Lys Pro Asn Pro Gly  
 195 200 205

Ser Ser Thr Asp Gly Asn Ser Ala Gly His Ser Gly Asn Asn Leu Leu  
 210 215 220

Gly Ser Glu Val Ala  
 225

- 16 -

<210> 7  
 <211> 687  
 <212> DNA  
 <213> Homo sapiens  
 <220>

<221> CDS  
 <222> (1)..(687)

<400> 7  
 atg gct gtg aga agg gac tcc gtg tgg aag tac tgc tgg ggt gtt ttg 48  
 Met Ala Val Arg Arg Asp Ser Val Trp Lys Tyr Cys Trp Gly Val Leu  
 1 5 10 15  
 atg gtt tta tgc aga act gcg att tcc aaa tcg ata gtt tta gag cct 96  
 Met Val Leu Cys Arg Thr Ala Ile Ser Lys Ser Ile Val Leu Glu Pro  
 20 25 30  
 atc tat tgg aat tcc tcg aac tcc aaa ttt cta cct gga caa gga ctg 144  
 Ile Tyr Trp Asn Ser Ser Asn Ser Lys Phe Leu Pro Gly Gln Gly Leu  
 35 40 45  
 gta cta tac cca cag ata gga gac aaa ttg gat att att tgc ccc aaa 192  
 Val Leu Tyr Pro Gln Ile Gly Asp Lys Leu Asp Ile Ile Cys Pro Lys  
 50 55 60  
 gtg gac tct aaa act gtt ggc cag tat gaa tat tat aaa gtt tat atg 240  
 Val Asp Ser Lys Thr Val Gly Gln Tyr Glu Tyr Tyr Lys Val Tyr Met  
 65 70 75 80  
 gtt gat aaa gac caa gca gac aga tgc act att aag aag gaa aat acc 288  
 Val Asp Lys Asp Gln Ala Asp Arg Cys Thr Ile Lys Lys Glu Asn Thr  
 85 90 95  
 cct ctc ctc aac tgt gcc aaa cca gac caa gat atc aaa ttc acc atc 336  
 Pro Leu Leu Asn Cys Ala Lys Pro Asp Gln Asp Ile Lys Phe Thr Ile  
 100 105 110  
 aag ttt caa gaa ttc agc cct aac ctc tgg ggt cta gaa ttt cag aag 384  
 Lys Phe Gln Glu Phe Ser Pro Asn Leu Trp Gly Leu Glu Phe Gln Lys  
 115 120 125

- 17 -

aac aaa gat tat tac att ata tct aca tca aat ggg tct ttg gag ggc 432  
 Asn Lys Asp Tyr Tyr Ile Ile Ser Thr Ser Asn Gly Ser Leu Glu Gly  
 130 135 140

ctg gat aac cag gag gga ggg gtg tgc cag aca aga gcc atg aag atc 480  
 Leu Asp Asn Gln Glu Gly Gly Val Cys Gln Thr Arg Ala Met Lys Ile  
 145 150 155 160

ctc atg aaa gtt gga caa gat gca agt tct gct gga tca acc agg aat 528  
 Leu Met Lys Val Gly Gln Asp Ala Ser Ser Ala Gly Ser Thr Arg Asn  
 165 170 175

aaa gat cca aca aga cgt cca gaa cta gaa gct ggt aca aat gga aga 576  
 Lys Asp Pro Thr Arg Arg Pro Glu Leu Glu Ala Gly Thr Asn Gly Arg  
 180 185 190

agt tcg aca aca agt ccc ttt gta aaa cca aat cca ggt tct agc aca 624  
 Ser Ser Thr Thr Ser Pro Phe Val Lys Pro Asn Pro Gly Ser Ser Thr  
 195 200 205

gac ggc aac agc gcc gga cat tcg ggg aac aac atc ctc ggt tcc gaa 672  
 Asp Gly Asn Ser Ala Gly His Ser Gly Asn Asn Ile Leu Gly Ser Glu  
 210 215 220

gtg gcc tta ttt gca 687  
 Val Ala Leu Phe Ala  
 225

<210> 8  
 <211> 229  
 <212> PRT  
 <213> Homo sapiens

<400> 8  
 Met Ala Val Arg Arg Asp Ser Val Trp Lys Tyr Cys Trp Gly Val Leu  
 1 5 10 15

Met Val Leu Cys Arg Thr Ala Ile Ser Lys Ser Ile Val Leu Glu Pro  
 20 25 30

Ile Tyr Trp Asn Ser Ser Asn Ser Lys Phe Leu Pro Gly Gln Gly Leu  
 35 40 45

- 18 -

Val Leu Tyr Pro Gln Ile Gly Asp Lys Leu Asp Ile Ile Cys Pro Lys  
 50 55 60

Val Asp Ser Lys Thr Val Gly Gln Tyr Glu Tyr Tyr Lys Val Tyr Met  
 65 70 75 80

Val Asp Lys Asp Gln Ala Asp Arg Cys Thr Ile Lys Lys Glu Asn Thr  
 85 90 95

Pro Leu Leu Asn Cys Ala Lys Pro Asp Gln Asp Ile Lys Phe Thr Ile  
 100 105 110

Lys Phe Gln Glu Phe Ser Pro Asn Leu Trp Gly Leu Glu Phe Gln Lys  
 115 120 125

Asn Lys Asp Tyr Tyr Ile Ile Ser Thr Ser Asn Gly Ser Leu Glu Gly  
 130 135 140

Leu Asp Asn Gln Glu Gly Gly Val Cys Gln Thr Arg Ala Met Lys Ile  
 145 150 155 160

Leu Met Lys Val Gly Gln Asp Ala Ser Ser Ala Gly Ser Thr Arg Asn  
 165 170 175

Lys Asp Pro Thr Arg Arg Pro Glu Leu Glu Ala Gly Thr Asn Gly Arg  
 180 185 190

Ser Ser Thr Thr Ser Pro Phe Val Lys Pro Asn Pro Gly Ser Ser Thr  
 195 200 205

Asp Gly Asn Ser Ala Gly His Ser Gly Asn Asn Ile Leu Gly Ser Glu  
 210 215 220

Val Ala Leu Phe Ala  
 225

## INTERNATIONAL SEARCH REPORT

In national Application No  
PCT/EP 01/11252

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07K14/715 C07K14/52 C12N5/10 A61K48/00 A61K38/19

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K C12N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, MEDLINE; BIOSIS, EMBASE, CHEM ABS Data, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5 670 625 A (BECKMANN M PATRICIA ET AL) 23 September 1997 (1997-09-23)  see whole doc. esp. examples, claims and col. 31.9ff, col 9 1.34 ff. ---	4, 7, 10, 11, 14, 19, 20, 23
X	WO 96 26958 A (HARVARD COLLEGE) 6 September 1996 (1996-09-06)  see whole doc. esp. claims and p.5, 1.14, p.8, 1.13 ff. ---	4, 5, 7, 8, 10-12, 14, 19, 20, 23
X	WO 99 17796 A (BRISKIN MICHAEL J ; LEUKOSITE INC (US); ZOU LILY (US)) 15 April 1999 (1999-04-15) see whole doc. esp. claims, and e.g. p.21.18; p.4, 1.8-21, ---	4, 8, 10, 11, 14, 19, 20, 23
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\* & \* document member of the same patent family

Date of the actual completion of the international search

10 January 2002

Date of mailing of the international search report

18/01/2002

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Mueller, F

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 01/11252

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 00 30673 A (GENENTECH INC) 2 June 2000 (2000-06-02) see whole doc. esp. claims and p.10, l.24-39; p.16, l.21-28; p.22, l.36-p.28., l.12) -----	1-23
X	WO 96 02645 A (GENENTECH INC) 1 February 1996 (1996-02-01)  see whole doc. esp. claims and seq ids 2,4 and 6 -----	4,7,10, 11,14, 19,20,23
X	WO 96 01839 A (IMMUNEX CORP) 25 January 1996 (1996-01-25)  see whole doc. esp. claims and seq id 2 -----	4,6,7, 11,14, 19,20,23

## INTERNATIONAL SEARCH REPORT

In International Application No

PCT/EP 01/11252

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5670625	A	23-09-1997	US 5512457 A	30-04-1996
			US 5728813 A	17-03-1998
			US 5627267 A	06-05-1997
			AU 669960 B2	27-06-1996
			AU 5667594 A	08-06-1994
			CA 2148484 A1	26-05-1994
			EP 0669929 A1	06-09-1995
			JP 8503368 T	16-04-1996
			NZ 258697 A	26-03-1996
			WO 9411384 A1	26-05-1994
WO 9626958	A	06-09-1996	WO 9626958 A2	06-09-1996
WO 9917796	A	15-04-1999	AU 9594598 A	27-04-1999
			EP 1019080 A2	19-07-2000
			JP 2001518318 T	16-10-2001
			WO 9917796 A2	15-04-1999
WO 0030673	A	02-06-2000	AU 1739700 A	13-06-2000
			EP 1135153 A1	26-09-2001
			WO 0030673 A1	02-06-2000
WO 9602645	A	01-02-1996	AU 705107 B2	13-05-1999
			AU 3008995 A	16-02-1996
			CA 2194955 A1	01-02-1996
			EP 0773997 A2	21-05-1997
			JP 10501701 T	17-02-1998
			WO 9602645 A2	01-02-1996
			US 5624899 A	29-04-1997
			US 5864020 A	26-01-1999
WO 9601839	A	25-01-1996	US 6303769 B1	16-10-2001
			AU 684988 B2	08-01-1998
			AU 3094095 A	09-02-1996
			CA 2194299 A1	25-01-1996
			EP 0770088 A1	02-05-1997
			FI 965297 A	07-03-1997
			JP 10502810 T	17-03-1998
			NO 965627 A	07-03-1997
			NZ 290253 A	26-02-1998
			WO 9601839 A1	25-01-1996
			US 2001009768 A1	26-07-2001